PALM BEACH ORTHOPAEDIC INSTITUTE, P.A.

G. CLAY BAYNHAM, M.D. | MICHAEL COONEY, M.D. | MELISA ESTES, M.D. | VINCENT FOWBLE, M.D. | KENNETH GERSZBERG, M.D. BRIAN HILL, M.D. | JOHN HINSON, M.D. | JAMES KERPSACK, M.D. | JUSTIN KEARSE, M.D. | MICHAEL LEIGHTON, M.D. | FERNANDO MORA, M.D. ANDREW NOBLE, M.D. | SCOTT NORRIS, D.O. | JOHN SCHILERO, D.P.M. | ANDREW SELTZER, D.O. | GARY WEXLER, M.D.

Rehabilitation Protocol Frozen Shoulder Release

Phase I: Maintain ROM (0 to 6 weeks)

- May remove dressing and shower postop day # 3.
- Sutures are all underneath the skin and will dissolve on their own
- Ice or cold flow systems encouraged for the first week at a minimum: should be used 3-4 times per day.
- Sling should be used for comfort only.
- Pulleys or Continuous Passive Motion (CPM) machine to be used 3-5 times per day.
- Supplement exercise program 3 times per day:
 Immediate elbow, forearm and hand range of motion out of sling Pendulum exercises
 Aggressive PROM and capsular mobility in all planes
- Supervised PROM and capsular stretching 3 times per week.
- May start active scapular mobility exercises immediately.
- Initiate AROM when tolerated.

Phase II: Restore Function (>6 weeks)

- Initiate gentle rotator cuff strengthening (Theraband, dumbbells, etc)
- Continue scapular stabilizer strengthening.
- Avoid strengthening in positions of impingement.