PALM BEACH ORTHOPAEDIC INSTITUTE, P.A.

G. CLAY BAYNHAM, M.D. | MICHAEL COONEY, M.D. | MELISA ESTES, M.D. | VINCENT FOWBLE, M.D. | KENNETH GERSZBERG, M.D. BRIAN HILL, M.D. | JOHN HINSON, M.D. | JAMES KERPSACK, M.D. | JUSTIN KEARSE, M.D. | MICHAEL LEIGHTON, M.D. | FERNANDO MORA, M.D. ANDREW NOBLE, M.D. | SCOTT NORRIS, D.O. | JOHN SCHILERO, D.P.M. | ANDREW SELTZER, D.O. | GARY WEXLER, M.D.

Rehabilitation Protocol Anterior Capsular Shift (Open)

Phase I: Protect Repair (0 to 6 weeks after surgery)

- Patients may shower immediately over clear plastic, waterproof dressing
- Sutures are all underneath the skin and will dissolve on their own
- Ice or cold flow systems encouraged for the first week at a minimum: should be used 3-4 times per day.
- Sling should be in place when not performing exercises.
- Initiate exercise program 3 times per day:

Immediate elbow, forearm and hand range of motion out of sling

Pendulum exercises

Passive and active assistive ER at the side to 30, flexion to 130

- May start active scapular mobility exercises at 3 to 4 weeks Must keep the shoulder musculature relaxed.
- AVOID range of motion into abduction, ER >30 degrees or ACTIVE IR

Phase II: Progress ROM & Protect Repair (6 to 12 weeks after surgery)

- May discontinue sling.
- Lifting restriction of 5 pounds with the involved extremity.
- Advance active and passive ROM:

ER at the side and flexion to tolerance

Scapular plane elevation to 130

IR and extension to tolerance

- Initiate gentle rotator cuff strengthening, EXCEPT IR.
- Continue scapular stabilizer strengthening.
- Avoid combined abduction and ER ROM, active or passive.
- Avoid active or resistive IR.

<u>Phase III:</u> Full Function (greater than 3 months after surgery)

- Begin combined abduction and ER ROM and capsular mobility.
- Discontinue lifting restrictions.
- Begin subscapularis strengthening.
- Advance rotator cuff and scapular stabilizer strengthening.
- Initiate functional progression to sports specific activities at 4 months.